

E_{11} , generalised space-time and IIA string theory

Peter West

Department of Mathematics

King's College, London WC2R 2LS, UK

As advocated in hep-th/0307098 we construct the non-linear realisation of the semi-direct product of E_{11} and its first fundamental representation at lowest level from the IIA viewpoint. We find a theory that is $SO(10,10) \otimes GL(1)$ invariant and contains the fields of gravity, a two form and a dilaton but which depend on coordinates which belong to the vector representation of $SO(10,10)$. The resulting Lagrangian agrees that of recent work on the so called doubled field theory. However, the construction given in this paper is straightforward and systematic. It also reveals the relevant underlying symmetries and opens the way to include the Ramond-Ramond, and higher level, fields together with additional coordinates of the generalised space-time.

When the E_{11} symmetry was first conjectured space-time was encoded into the non-linear realisation by introducing the space-time translation generators in an ad hoc manner [1,2]. It was subsequently proposed [3] that one should introduce generators transforming in the fundamental representation l_1 of E_{11} ; more precisely one should take the non-linear realisation of the semi-direct product of E_{11} and a set of generators in the fundamental representation associated with node one (see figure 1) , i.e. $E_{11} \otimes_s l_1$ [3]. At lowest levels the l_1 multiplet in eleven dimensions begins with the space-time translation generators P_a , then a two form $Z^{a_1 a_2}$, a five form generator $Z^{a_1 \dots a_5}$ and a generator $Z^{a_1 \dots a_7, b}$ together with an infinite number of other generators. In this approach the fields in eleven dimensions would depend on all the coordinates introduced in this non-linear realisation that is $x^a, x_{a_1 a_2}, x_{a_1 \dots a_5}, x_{a_1 \dots a_7, b}, \dots$ [3].

There is convincing evidence that all the brane charges are contained in the l_1 representation [3,4,5,6,7] and the introduction of the different coordinates corresponds to measuring using the different brane probes. The non-linear realisation based on $E_{11} \otimes_s l_1$ is a theory that possess many new unfamiliar coordinates, indeed at all levels an infinite number of coordinates and it is not clear how to recover a field theory which has only the dependence on the usual space-time. Nonetheless the non-linear realisation of $E_{11} \otimes_s l_1$ has been used to construct the gauged supergravities with maximal supersymmetry, at least the field strengths of the four dimensional theories, although the techniques used are apply to all dimensions [8]. In this case not all the coordinates of the l_1 representation were used but instead of taking just the usual space-time coordinates x^a one took a space-time to be a slice which lies in l_1 and E_{11} and contains many new coordinates.

A new generalised geometry was proposed in references [8,9] and then subsequent to reference [3] used in a large number of papers to reformulate parts of supergravity theories. However, reference [3] automatically introduces a generalise tangent space with a generalised vierbein which is easily computed and whose role in the theory is very strongly constrained by the symmetries of the non-linear realisation. It is clear that many features of the work using generalised geometry are automatically encoded in the non-linear realisation of $E_{11} \otimes_s l_1$.

In this paper we will compute the non-linear realisation of $E_{11} \otimes_s l_1$ at the lowest level appropriate for the IIA theory. This has a $SO(10,10) \otimes GL(1)$ symmetry and the same fields as the NS-NS sector of the superstring, however, these fields depend on the coordinates x^a, y_a that belong to the vector representation of $SO(10,10)$. Following the ideas put forward in [11,1,2,12] we consider the fields to only depend on x^a and demand that the theory is invariant under diffeomorphisms. This fixes the coefficients in the Lagrangian and we arrive at the Lagrangian for the Neveu-Schwarz-Neveu-Schwarz sector of IIA supergravity. A key role is played by the $GL(1)$ part of the symmetry without which a group theoretic derivation of the result would not have been possible. The coordinates x^a, y_a were first introduced in the context of string theory in reference [13,14]. In a recent paper [15] the the non-linear realisation of $E_{11} \otimes_s l_1$ at lowest level appropriate for the IIA theory was used to deduce a formulation [13] of the bosonic sector of the superstring.

To find a ten dimensional theory from the E_{11} non-linear realisation one must select an A_9 , or $Sl(10)$, subalgebra, the so called gravity line, as this subalgebra leads to ten dimensional gravity. Looking at the Dynkin diagram of E_{11} , see figure 1, one sees that

there are only two possibilities; the nodes from one to nine inclusive and the nodes one to eight inclusive and node eleven. The latter leads to the IIB theory [16] and the former the IIA theory [2] which is the subject of this paper. There are two nodes in the IIA theory which are not associated with gravity, that is nodes ten and eleven. Deleting node ten (see fig 1) leaves the algebra D_{10} , in particular its real form $SO(10,10)$, and it is useful to decompose the E_{11} adjoint representation, which is the one that occurs in the non-linear realisation, into representations of $SO(10,10)$. However, to recognise objects that are more familiar it is helpful to further decompose these representations into those of $SL(10)$ which corresponds to deleting node eleven in addition.

The representations that occur in a decomposition of the adjoint representation of E_{11} to a subalgebra associated with a deleted node can be classified in terms of increasing levels, whose precise definition is given in appendix A. The decomposition into representations of $SO(10,10)$, associated with the deletion of node ten, at level zero is just the adjoint representation of $SO(10,10)$ together with one other generator which is in the Cartan subalgebra of E_{11} . When written in terms of representations of $SL(10)$ the generators of the adjoint representation of $SO(10,10)$ are $K^a{}_b$, R^{ab} , $\tilde{R}_{ab}$, $a, b = 1, 2, \dots, 10$, where the $K^a{}_b$ are the generators of the adjoint representation of the $SL(10)$ of the chosen gravity line. Their other generator we will denote as $\tilde{R}$. Their algebra is derived from E_{11} in appendix A and is given by

$$[K^a{}_b, K^c{}_d] = \delta_b^c K^a{}_d - \delta_d^a K^c{}_b, \quad [K^a{}_b, R^{cd}] = \delta_b^c R^{ad} - \delta_b^d R^{ac}, \quad [K^a{}_b, \tilde{R}_{cd}] = -\delta_c^a \tilde{R}_{bd} + \delta_d^a \tilde{R}_{bc},$$

$$[R^{ab}, \tilde{R}_{cd}] = 4\delta_{[c}^{[a} K^{b]}_{d]} + \frac{2}{3}\tilde{R}, \quad [R^{ab}, R^{cd}] = 0 = [\tilde{R}_{ab}, \tilde{R}_{cd}], \quad [\tilde{R}, R^{cd}] = 0, \quad [\tilde{R}, \tilde{R}_{cd}] = 0 \quad (1)$$

where $\tilde{R} = -\sum_{a=1}^{11} K^a{}_a + 3K^{11}{}_{11}$. We will show in appendix A that this algebra is that of $SO(10,10) \otimes GL(1)$ where the last factor has the generator $\tilde{R}$.

In the non-linear realisation of E_{11} the generators of the Borel subgroup lead to fields and so at level zero we find the fields $h_a{}^b$, $A_{a_1 a_2}$ and a that is the fields of the NS-NS sector of the IIA string. At the next level, i.e. level one, one finds generators, and so fields, which are all anti-symmetric tensors of $SL(10)$ which have odd ranks, i.e. those of one to nine and these belong to the Majorana Weyl spinor representation of $SO(10,10)$. These include the fields of the Ramond-Ramond sector and their duals as well as a rank nine form field that gives rise to the massive IIA theory. At higher levels one finds infinite number of fields many of which have a complicated index structure. In fact the six form field which is the dual of the two form field at level zero occurs at level two.

As explained in reference [3] to introduce a generalised space-time into the non-linear realisation we must also consider the fundamental representation of E_{11} associated with node one, denoted by l_1 , see appendix A. In particular we will consider the non-linear realisation of the semi-direct product of E_{11} and a set of generators that transform under E_{11} transformations, i.e E_{11} commutators, like the l_1 representation; we denote this semi direct product as $E_{11} \otimes_s l_1$. The coordinates of the generalised space-time arise as the parameters of the l_1 generators in the general group element. At level zero with respect to node ten the generators of the l_1 representation are $l_1^0 = \{P_a, Q^a, a = 1, 2, \dots, 10\}$ which belong to the vector representation of $SO(10,10)$. Their commutation relations with those

of $SO(10, 1) \otimes GL(1)$ are derived in appendix A from the $E_{11} \otimes_s l_1$ algebra and are given by

$$\begin{aligned}
[K^c_b, P_a] &= -\delta_a^c P_b + \frac{1}{2} \delta_b^c P_a, \quad [R^{ab}, P_c] = -\frac{1}{2} (\delta_c^a Q^b - \delta_c^b Q^a) = -\delta_c^{[a} Q^{b]}, \quad [\tilde{R}_{ab}, P_c] = 0, \\
[K^a_b, Q^c] &= \delta_b^c Q^a + \frac{1}{2} \delta_b^c Q^a \quad [\tilde{R}_{ab}, Q^c] = 4(\delta_a^c P_b - \delta_b^c P_a) = 2\delta_{[a}^c P_{b]}, \quad [R^{ab}, Q^c] = 0, \\
[\tilde{R}, P_a] &= -3P_a, \quad [\tilde{R}, Q^a] = -3Q^a
\end{aligned} \tag{2}$$

At level one the generators of l_1 , and so the associated coordinates, are all tensors of $SL(10)$ of even rank and they belong to the Majorana Weyl spinor representation of $SO(10, 10)$. It is of opposite chirality to those of the level one fields mentioned above.

We can now construct the non-linear realisation of $E_{11} \otimes_s l_1$ with local sub-algebra $K(E_{11})$ at level zero where $K(G)$ denotes the Cartan involution invariant subalgebra of the algebra G . This is just the non-linear realisation of $(E_{11} \otimes_s l_1)^{(0)} = (SO(10, 10) \otimes GL(1)) \otimes_s l_1^0$, whose algebra is given in equations (1) and (2) above. The local subalgebra of the non-linear realisation is just the Cartan involution invariant subalgebra of $SO(10, 10)$ which is $SO(10) \otimes SO(10)$. The non-linear realisation is built from a general element of $(E_{11} \otimes_s l_1)^{(0)}$ which is taken to transform as $g \rightarrow g_0 g h$ where the rigid transformation $g_0 \in (E_{11} \otimes_s l_1)^{(0)}$ and the local transformation $h \in SO(10) \otimes SO(10)$. Using the latter we can bring the general group element of $(E_{11} \otimes_s l_1)^{(0)}$ to be of the form

$$g = g_l g_E, \quad \text{where} \quad g_l = e^{x^a P_a + y_a Q^a} \quad \text{and} \quad g_E = e^{h \cdot K} e^{\frac{1}{2} A \cdot R} e^{a R} \tag{3}$$

where $R = \frac{1}{12}(9K^{11}_{11} - \sum_{a=1}^{11} K^a_a)$ and the fields $h_a^b, A_{a_1 a_2}$ and a depend on x^a and y_a . We note that we are using a different scalar generator R rather than the $\tilde{R}$ that appeared in equation (1) and it has the commutators

$$[R, P_a] = 0, \quad [R, Q^a] = \frac{1}{2} Q^a, \quad [R, K^a_b] = 0, \quad [R, R^{ab}] = \frac{1}{2} R^{ab}, \quad [R, R_{ab}] = -\frac{1}{2} R_{ab} \tag{4}$$

The Cartan form is given by

$$\mathcal{V} = g^{-1} dg = g_E^{-1} (dx^a P_a + dy_a Q^a) g_E + g_E^{-1} dg_E \tag{5}$$

The first term is given by

$$g_E^{-1} (dx^a P_a + dy_a Q^a) g_E \equiv (dx^a, dy_a) \mathcal{E} \begin{pmatrix} P_b \\ Q^b \end{pmatrix} = dz^T \mathcal{E} L \tag{6}$$

where the $2D$ by $2D$ matrix $\mathcal{E}$ is given by

$$\mathcal{E} = (dete)^{-\frac{1}{2}} \begin{pmatrix} e & \frac{1}{2} e A e^{-\frac{1}{2} a} \\ 0 & e^{-1T} e^{-\frac{1}{2} a} \end{pmatrix} \tag{7}$$

where $e = e_\mu^a = (e^h)_\mu^a$ and A is the matrix $A_{a_1 a_2}$. We can think of $\mathcal{E}$ as a generalised vielbein. In the last line we have use the definitions $dz^T = (dx^a, dy_a)$ and $L = \begin{pmatrix} P_b \\ Q^b \end{pmatrix}$. We note that $\mathcal{E}$ does not have determinant one, but this is consistent with the presence of the additional $GL(1)$ generator in the algebra whose non-linear realisation we are constructing.

The Cartan form is inert under the above rigid transformations but transforms under the local transformation h as $\mathcal{V} \rightarrow h^{-1}\mathcal{V}h + h^{-1}dh$. However, carrying out a rigid transformation $g \rightarrow g_0 g$ on the general group element of equation (3) we find that $g_l \rightarrow g_0 g_l g_0^{-1}$ and $g_E \rightarrow g_0 g_E$ as the l_1 generators form a representation of E_{11} . As a result the coordinates form a representation of E_{11} , and in particular at level zero x^a and y_a form a representation of $E_{11}^{(0)} \equiv SO(10,10) \otimes GL(1)$ and the action of rigid $E_{11}^{(0)}$ transformations on the coordinates is given by

$$x^a P_a + y_a Q^a \rightarrow x'^a P_a + y'_a Q^a = g_0(x^a P_a + y_a Q^a)g_0^{-1} \quad (8)$$

As expected they transform according to the vector representation of $SO(10,10)$.

We can define the action of the vector representation of $k \in E_{11}^{(0)} = SO(10,10) \otimes GL(1)$ using the generators of $l_1^{(0)}$ by $U(k)(L_N) \equiv k^{-1}L_N k = D(k)_N^M L_M$ where $L_N = \begin{pmatrix} P_b \\ Q^b \end{pmatrix}$ and $D(k)_N^M$ is the matrix representative. As a result of equation (8), we find that $dz^T \rightarrow dz^{T'} = dz^T D(g_0^{-1})$ or putting in the indices $dz^{TN} \rightarrow dz^{TN'} = dz^{TM} D(g_0^{-1})_M^N$ where as before $dz^{TN} = (dx^a, dy_a)$. The derivative $\partial_N = \frac{\partial}{\partial z^N}$ in the generalised space-time transforms as $\partial'_N = D(g_0)_N^M \partial_M$. Examining equation (5), we note that $\mathcal{E}$ is equal to the matrix $D(g_E)_N^M$. Almost identical considerations hold at all levels in the $E_{11} \otimes_s l_1$ non-linear realisation.

As the Cartan form is inert under rigid transformations, their action on the coordinates must be compensated by that on $\mathcal{E}$ which we give the indices $\mathcal{E}_N^A$ and so $\mathcal{E}_N^{A'} = D(g_0)_N^M \mathcal{E}_M^A$. The generalised vielbein $\mathcal{E}_N^{A'}$ transforms on its upper index by a local $SO(10) \otimes SO(10)$ transformation and so we can think of the upper index as a tangent index and the lower index as a world index.

Rather than use the Cartan forms we will construct the action out of $M \equiv g_E I_c(g_E^{-1})$ where I_c is the Cartan involution. It is easy to see that M is inert under local transformations but transforms as $M \rightarrow M' = g_0 M I_c(g_0^{-1})$ under rigid transformations. Fortunately at level zero, that is for the group $E_{11}^{(0)}$, $I_c(g^{-1}) = g^T$. It follows from the above discussion just above, and equation (7) that in the vector representation M takes the form

$$D(M) = \mathcal{E}\mathcal{E}^T = (dete)^{-1} \begin{pmatrix} ee^T - \frac{1}{4}eAAe^T e^{-a} & \frac{1}{2}eAe^{-1}e^{-a} \\ -\frac{1}{2}e^{-1T}Ae^T e^{-a} & e^{-1T}e^{-1}e^{-a} \end{pmatrix} \quad (9)$$

Writing out the indices explicitly $D(M)_{NM} = (\mathcal{E}\mathcal{E}^T)_{NM}$ it is clear that $D(M)$ is inert under local $SO(10) \otimes SO(10)$ transformations as the tangent index is summed but transforms as a $SO(10,10)$ vector on both of its lower world indices. In what follows we write $D(M)$ just as M for simplicity.

We can construct an invariant Lagrangian out of M and ∂_N . These are inert under local transformation and so the most general such object bilinear in generalised space-time

derivatives is given by

$$\begin{aligned}
L = & c_1 \partial_S M_{PQ} \partial_T (M^{-1})^{PQ} (M^{-1})^{ST} + c_2 \partial_S (M^{-1})^{PQ} \partial_P M_{TQ} (M^{-1})^{ST} \\
& + c_3 \partial_S (M^{-1})^{SR} \partial_P (M^{-1})^{PQ} M_{RQ} + \frac{c_4}{400} \partial_S \det M \partial_T (\det M)^{-1} (M^{-1})^{ST} \\
& + \frac{c_5}{10} \det M \partial_T (\det M)^{-1} \partial_S (M^{-1})^{ST}
\end{aligned} \tag{10}$$

where $c_1, \dots, c_5$ are constants.

It is instructive to evaluate this Lagrangian in terms of the fields and this is easiest achieved by going to string frame. That is carrying out the field redefinitions $e \rightarrow e^{-\frac{1}{4}a} e_s, A \rightarrow e^{\frac{1}{2}a} A_s$ whereupon we find that

$$\mathcal{E} = e^{-\frac{\tau}{2}} \tilde{\mathcal{E}}, \quad \text{where} \quad \tilde{\mathcal{E}} = \begin{pmatrix} e_s & \frac{1}{2} e_s A_s \\ 0 & e_s^{-1T} \end{pmatrix} \tag{11}$$

and $e^\tau = e^{-2a} \det e_s$. The Lagrangian of equation (10) then becomes

$$\begin{aligned}
L = & e^\tau (c_1 \partial_S \tilde{M}_{PQ} \partial_T (\tilde{M}^{-1})^{PQ} (\tilde{M}^{-1})^{ST} + c_2 \partial_S (\tilde{M}^{-1})^{PQ} \partial_P \tilde{M}_{TQ} (\tilde{M}^{-1})^{ST} \\
& + c_3 \partial_S (\tilde{M}^{-1})^{SR} \partial_P (\tilde{M}^{-1})^{PQ} \tilde{M}_{RQ}) - (20c_1 + c_2 - c_3 + c_4 - 2c_5) e^\tau \partial_S \tau \partial_T \tau (\tilde{M}^{-1})^{ST} \\
& - 2e^\tau (c_2 - c_3 - c_5) \partial_S \tau \partial_T (\tilde{M}^{-1})^{ST}
\end{aligned} \tag{12}$$

where $\tilde{M} = \tilde{\mathcal{E}} \tilde{\mathcal{E}}^T$. We note that although $\mathcal{E}$ does not have determinant one, $\tilde{\mathcal{E}}$ does have determinant one and is an element of $\text{SO}(10,10)$.

The field redefinition could also have been achieved by using a different choice of generators in the group element. Indeed, we can rewrite the group element of equation (3) in terms of the generators of $\text{SO}(10,10)$, given in appendix A, and $\tilde{R}$ to find

$$g_E = e^{h_s \cdot K} e^{\frac{1}{2} A_s \cdot R} e^{\frac{1}{3} a \tilde{R}} = e^{h_s \cdot \tilde{K}} e^{\frac{1}{2} A_s \cdot R} e^{\frac{1}{3} \tilde{a} \tilde{R}} \tag{13}$$

where $e_s = e^{h_s}$, as a matrix, and $e^{-2\tilde{a}} = e^\tau$. Thus string frame is associated with the group element when written in terms of the $\text{SO}(10,10)$ generators and e^τ is associated with the $\text{GL}(1)$ factor.

So far we have constructed a theory which is invariant under a rigid symmetry namely $\text{SO}(10,10) \otimes \text{GL}(1)$ but we would like to construct a theory that has local symmetries which replace the rigid symmetries of $\text{SO}(10,10)$. This is a theory with diffeomorphisms and gauge symmetries. The way to do this has been set out in the context of other theories. For the case of gravity one writes down the non-linear realisation of $\text{GL}(D) \otimes_s P$ where P are the translations and then demands that the theory admit diffeomorphism, or equivalently, admit a simultaneous realisation with the conformal group [11]. This fixes the constants in the Lagrangian which has just the rigid symmetries of the non-linear realisation and one finds Einstein's theory of gravity. A similar procedure was followed at low levels for the $E_{11} \otimes l_1$ non-linear realisation to find maximal supergravity theories but in these theories one also finds that the theory was gauge invariant corresponding to the

presence of the gauge fields [1,2]. A more subtle strategy was carried out in the context of maximal supergravity in four dimensions and its E_7 symmetry [12]. An $E_7 \otimes_s l_1$ non-linear realisation was constructed where in this case the l_1 representations is the fifty six dimensional representation of E_7 . It was shown [12] that the corresponding action admitted a diffeomorphism symmetry if one neglected the coordinate dependence on forty nine of the fifty six coordinates, leaving a seven dimensional dependence, and fixed the constants in the action to a particular set of values. In fact this diffeomorphism invariance required a contribution from the four dimensional metric of the eleven dimensional supergravity theory in which the E_7 theory was embedded. Unlike the case of gravity, the E_7 theory can not admit a conformal symmetry that would imply the rigid symmetries of the non-linear realisation become local.

We now follow this same strategy here. We restrict the derivatives in the generalised space-time to be only in the x^a directions. Carrying out this step let us evaluate the Lagrangian of equations (10) and (12) by substituting the expression for $\tilde{\mathcal{E}}$ of equation (11), we find that

$$\begin{aligned}
L = & c_1 e^\tau (2\partial_\mu g_{\rho\kappa} \partial^\mu g^{\rho\kappa} - \frac{1}{2} \partial_\mu A_\rho{}^\kappa \partial^\mu A^\rho{}_\kappa - \frac{1}{2} \partial_\mu A_{\rho\kappa}^2 \partial^\mu g^{\rho\kappa}) \\
& + c_2 e^\tau (g^{\rho\mu} \partial_\mu g^{\nu\kappa} \partial_\nu g_{\rho\kappa} - \frac{1}{4} \partial^\rho A^\nu{}_\kappa \partial_\nu A_\rho{}^\kappa + \frac{1}{4} \partial_\mu g^{\nu\kappa} \partial_\nu A^\rho{}_\kappa A^\mu{}_\rho + \frac{1}{4} \partial_\nu g^{\rho\kappa} \partial_\mu A^\nu{}_\kappa A^\mu{}_\rho - \frac{1}{4} \partial^\rho g^{\nu\kappa} \partial_\nu A_{\rho\kappa}^2) \\
& + c_3 e^\tau (\partial_\mu g^{\mu\rho} \partial_\nu g^{\nu\kappa} g_{\rho\kappa} + \frac{1}{4} \partial_\mu A^\mu{}_\rho \partial_\nu A^\nu{}_\kappa g^{\rho\kappa} + \frac{1}{2} \partial_\mu A^\mu{}_\rho \partial_\nu g^{\nu\kappa} A^\rho{}_\kappa - \frac{1}{4} \partial_\mu g^{\mu\rho} \partial_\nu g^{\nu\kappa} A_{\rho\kappa}^2) \\
& - (20c_1 + c_2 - c_3 + c_4 - 2c_5) e^\tau \partial_\mu \tau \partial_\nu \tau g^{\mu\nu} - 2e^\tau (c_2 - c_3 - c_5) \partial_\mu \tau \partial_\nu g^{\mu\nu} \quad (14)
\end{aligned}$$

where $\partial^\mu = g^{\mu\nu} \partial_\nu$ and $A_{\mu\nu}^2 = A_\mu{}^\rho A_{\rho\nu}$.

To fix the coefficients in the Lagrangian it is simplest to carry out an infinitesimal diffeomorphism and gauge transformation on the bilinear terms and in particular the variations $\delta h_{\mu\nu} = \partial_\mu \xi_\nu + \partial_\nu \xi_\mu$, where $g_{\mu\nu} = e_\mu{}^a e_{\nu a} = \eta_{\mu\nu} + 2h_{\mu\nu}$ at lowest order, and $\delta A_{\mu\nu} = \partial_\mu \Lambda_\nu - \partial_\nu \Lambda_\mu$ we find the action is invariant provided $c_1 = -\frac{c_2}{4}$, $c_4 = 6c_2$, $c_5 = 0$. In fact the coefficient c_2 and c_3 occur in the combination $c_2 - c_3$ and so we can set $c_3 = 0$. We can choose the overall scale of the Lagrangian so that

$$c_1 = \frac{1}{4}, c_2 = -1, c_4 = -6 \quad (15)$$

all others being zero.

We find that with these constants the Lagrangian is just that for the NS-NS sector of the IIA supergravity which is given by

$$4 \int d^{10}x \det e e^{-2a} \left\{ \frac{1}{2} R - \frac{1}{3.8.4} F_{\mu_1 \mu_2 \mu_3} F^{\mu_1 \mu_2 \mu_3} + 2\partial_\mu a \partial^\mu a \right\} \quad (16)$$

In deriving this result we have used the identity

$$\int d^D x \sqrt{-\det g} e^{-2a} R$$

$$\begin{aligned}
&= \int d^D x \sqrt{-\det g} e^{-2a} \left\{ -\frac{1}{2} \partial^\tau g^{\nu\lambda} \partial_\nu g_{\tau\lambda} + \frac{1}{4} \partial_\nu g_{\rho\kappa} \partial^\nu g^{\rho\kappa} + \frac{1}{4} \partial_\nu (\ln \det g) \partial^\nu (\ln \det g) \right. \\
&\quad \left. - \frac{1}{2} \partial^\nu (\ln \det g) \partial^\mu g_{\mu\nu} - 2 \partial_\nu a (2\Gamma^\nu - g^{\nu\tau} \partial^\rho g_{\tau\rho}) \right\} \\
&= \int d^D x \sqrt{-\det g} e^{-2a} \left\{ -\frac{1}{2} \partial^\tau g^{\nu\lambda} \partial_\nu g_{\tau\lambda} + \frac{1}{4} \partial_\nu g_{\rho\kappa} \partial^\nu g^{\rho\kappa} + \partial_\mu \tau \partial^\mu \tau - \partial^\mu \tau \partial^\lambda g_{\mu\lambda} - 4 \partial^\mu a \partial_\mu a \right\}
\end{aligned} \tag{17}$$

valid in D dimensions and the definitions $\Gamma_{\mu\nu}^\lambda = \frac{1}{2} g^{\lambda\tau} (\partial_\nu g_{\tau\mu} + \partial_\mu g_{\tau\nu} - \partial_\tau g_{\mu\nu})$ and $R_{\mu\nu}{}^\rho{}_\lambda = \partial_\mu \Gamma_{\nu\lambda}^\rho + \Gamma_{\mu\kappa}^\rho \Gamma_{\nu\lambda}^\kappa - (\mu \rightarrow \nu)$.

It is interesting to trace how the familiar density factor $\det e$ arises in the above Lagrangians. Each factor of M carries with it a $(\det e)^{-1}$ factor which it inherited from the product of two $\mathcal{E}$'s each with factor $(\det e)^{-\frac{1}{2}}$, this in turn had its origin in the last terms, with factor of one half, in the commutation relations of K^a_b with P_a and Q^a given in equation (2). This additional term arises as P_a is the highest weight state in the E_{11} representation. The two space-time derivatives in the Lagrangian ensure that there is one more factor of M^{-1} than of M in order to balance the indices and as a consequence we find the desired $\det e$ factor. It is interesting to note that this factor has its origins in the E_{11} algebra and the definition of the l_1 representation.

We now comment on the relation of this paper to the work on the so called double field theory. This result was developed in a number of substantial papers [17-20] which had their origins in earlier work on string field theory [21] and reference [22]. The final result was a field theory with a metric, two form and dilaton, i.e. the fields of the massless NS-NS sector of closed strings, but which depend on the coordinates x^a and y_a which transformed as a vector of $O(D,D)$. As the author of this paper understands it, doubled field theory was found by a circuitous route beginning [17] by extracting the quadratic and cubic terms, together with their gauge transformations, from gauge covariant closed string field theory [23]. These terms were then shown to be invariant under a set of $O(10,10)$ transformations introduced by ansatz. By assuming a set of requirements, gauge transformations were found to all orders in the fields and written in an $O(10,10)$ covariant way [18]. By introducing covariant derivatives an all orders Lagrangian was found that agreed with that of closed string field theory at quadratic and cubic order [19] and shown to be invariant under the gauge transformations if a constraint held. In the general case this constraint was the same as suppressing all dependence on y_a .

Finally, this Lagrangian was expressed in terms of an $O(10,10)$ generalised metric H (the $\tilde{M}$ in this paper) and e^{-2d} (the $e^{-\tau}$ in this paper). In fact this Lagrangian was previously contained in a seminar [24] of one of the authors (OH). However, it was also claimed in the seminar that $H^{-1} \partial H$ was the Cartan form of $O(10,10)$. It was pointed out [25] by the author of this paper that the final result was almost certainly the lowest level non-linear realisation of $E_{11} \otimes_s l_1$, i.e. the result of this paper, and that the non-linear realisation would contain $\tilde{M} = gg^T$ where $g \in O(10,10)$ with the transformation $g \rightarrow g_0 g h$ being understood. This relation subsequently appeared in paper [20] but in the form $\tilde{M} = g^T g$. Unfortunately this latter expression is not inert under the local h part to the transformation $g \rightarrow g_0 g h$ of equation (5.1) of paper [20].

Although our final result of equation (10) with the constants of equation (15) agrees with that of [20] the derivation presented in this paper is very different. We begin with an entirely group theoretic construction, i.e the non-linear realisation of $E_{11} \otimes_s l_1$, as proposed in reference [3], at lowest level. It is important to realise that this is not the same as the traditional sigma model associated with internal symmetries. The fields, their rigid $O(D, D) \otimes GL(1)$ transformations and the most general Lagrangian are found by a very straightforward calculation. The constants in the Lagrangian are fixed, following the ideas of [11,1,2,12], by demanding that the Lagrangian be diffeomorphism invariant when the fields are restricted to depend only on x^μ . Crucial to this construction is the $GL(1)$ factor in the symmetry which seems to play no role in the papers on double field theory. Such an additional factor is required from the non-linear realisation perspective since the generators not in the local subalgebra lead to fields in the final theory and we require a $D + 1$ commuting subalgebra in order to find the diagonal components of the metric and the dilaton. As explained above it is also essential to get the usual $\det e$ density factor in the Lagrangian.

The $E_{11} \otimes_s l_1$ non-linear realisation also provides a clear method to include the Ramond-Ramond fields and indeed higher level fields and also the corresponding additional coordinates. Furthermore it makes it clear that the symmetry underlying the doubled field theory, which had its origins in string field theory, is just that of the non-linear realisation of $E_{11} \otimes_s l_1$.

It is instructive to consider the eleven dimensional theory formed from the $E_{11} \otimes_s l_1$ non-linear realisation with local subalgebra $K(E_{11})$ at lowest level, i.e. level zero with respect to the deletion of node eleven. The resulting subalgebra is $GL(11)$ which is associated with eleven dimensional gravity. At level zero one has the $GL(11)$ algebra, with generators K^a_b , $a, b = 1, \dots, 11$ and the local subalgebra is just $SO(10)$, while in the l_1 representation there are only the usual space-time translations P_a . Their commutation relations, as derived from E_{11} , are given by

$$[K^a_b, K^c_d] = \delta_b^c K^a_d - \delta_d^a K^c_b, \quad [K^c_b, P_a] = -\delta_a^c P_b + \frac{1}{2} \delta_b^c P_a \quad (18)$$

We note the presence of the second term in the last commutator whose origin was discussed in appendix A. We can now follow exactly the path as above, we take the group element $g = g_l g_E = e^{x \cdot P} e^{h \cdot K}$ and find from the Cartan form that $\mathcal{E}_\mu^a = (\det e)^{-\frac{1}{2}} e$ where $e_\mu^a = (e^h)_\mu^a$. The most general invariant action is formed from $\partial_\mu = \frac{\partial}{\partial x^\mu}$ and $M = \mathcal{E} \mathcal{E}^T$ and has the above form of equation (10) with the indices $M, N, \dots$ now being replaced by $\mu, \nu = 1, \dots, 11$. We find a diffeomorphism invariant result, which is general relativity, in D dimensions provided $c_1 = \frac{1}{4}$, $c_2 = -\frac{1}{2}$, $c_3 = 0 = c_5$ and $c_4 = -\frac{1}{D-2}$. Of course $D = 11$ for the case of interest to us. However, we note that the $\det e$ factor has the same origin as that just described above, namely in the second term of the last commutator of equation (18).

It is amusing to consider the next level in eleven dimensions. At level one we have the three form generator R^{abc} in the adjoint representation of E_{11} and the generators Z_{ab} in l_1 representation. Thus we have the field A_{abc} which now depends on the coordinates x^a and y_{ab} . The group element is $g = g_l g_E$ with $g_l = e^{x \cdot P} e^{y \cdot Z}$ and $g_E = e^{h \cdot K} e^{A \cdot R}$. The

Cartan form takes the form

$$\begin{aligned}\mathcal{V} &= g^{-1}dg = g_E^{-1}(dx^a P_a + dy_{ab} Z^{ab})g_E + g_E^{-1}dg_E \\ &= (dx^a, dy_{ab})\mathcal{E} \begin{pmatrix} P_b \\ Z^{ab} \end{pmatrix} + g_E^{-1}dg_E\end{aligned}\tag{19}$$

where the generalised vierbein $\mathcal{E}$ is given as a matrix by

$$\mathcal{E} = (dete)^{-\frac{1}{2}} \begin{pmatrix} e_\mu^a & -\frac{1}{2}e_\mu^c A_{cb_1 b_2} \\ 0 & (e^{-1})_{[b_1}^{\mu_1} (e^{-1})_{b_2]}^{\mu_2} \end{pmatrix}\tag{20}$$

One can then construct $M = \mathcal{E}\mathcal{E}^T$ which is invariant under local transformations. We hope to report elsewhere on the dynamics of the non-linear realisation. The same calculation can easily be carried out for the IIA theory at the next level. The additional fields and coordinates were listed earlier in this paper.

In a previous paper [15] it was shown that quantising the $SO(10,10)$ symmetric string leads to a field theory with coordinates x^a , or y^a or a field theory with the coordinates x^a and y^a but which obey non-trivial commutation relations, that is a non-commutative field theory. The different theories corresponding to the different choices of representations of the fundamental commutators of the theory. It is tempting to assume that all these different theories lead to the same physical result, but it can not be excluded that they may differ in subtle ways. It would be interesting to investigate the connection with the results found in this paper. It is striking that some non-linear realisations admit, with a suitable choice of constants, local symmetries. The deeper meaning of this result has yet to be understood.

Acknowledgment

The author wishes to thank Christian Hillmann for extensive discussions that substantially helped the development of this work and the Erwin Schrödinger International Institute for Mathematical Physics, Vienna for providing financial support and hospitality during the author's stay in Vienna where much of this work was carried out. He also thanks the Physics department of the Technical University of Vienna for their hospitality and the STFC for support from the rolling grant awarded to King's.

Appendix A

In this appendix we calculate the algebra $E_{11} \otimes_s l_1$ at lowest level for the decomposition appropriate to the IIA theory. We will begin with the known algebra of the E_{11} algebra in terms of A_{10} , or $SL(11)$, representations [1,3]. This algebra is found by deleting node eleven in the Dynkin Diagram of E_{11} (see figure 1). For a decomposition of E_{11} corresponding to a subalgebra associated with the deletion of a particular node, the resulting generators can be classified in terms of increasing level. As for any Kac-Moody algebra, the generators of E_{11} are constructed as multiple commutators of the Chevalley generators and the level of a given positive (negative) root generator with respect to a particular node is just plus (minus) the number of times that the positive (negative) root Chevalley generators

associated with this node occurs in this multiple commutator. For the decomposition to SL(11) we consider the level associated with node eleven. At level zero we have the algebra GL(11) with the generators K^a_b , $a, b = 1, \dots, 11$ and at level one and minus one the rank three generators R^{abc} and R_{abc} . These level zero and one generators contain all the Chevalley generators of E_{11} (the positive root Chevalley generator associated with node eleven is $R^{9^{10}11}$) and so their multiple commutators lead to all generators of E_{11} . As a result the level of a generator that occurs in the SL(11) decomposition is just the number of times the generator R^{abc} occurs minus the number of times the generator R_{abc} occurs. The generators at level two and minus two are $R^{a_1 \dots a_6}$ and $R_{a_1 \dots a_6}$ respectively, while those at levels three and minus three are $R^{a_1 \dots a_8, b}$ and $R_{a_1 \dots a_8, b}$ respectively

The E_{11} algebra at levels zero and up three is given by [1,3]

$$[K^a_b, K^c_d] = \delta_b^c K^a_d - \delta_d^a K^c_b, \quad (A.1)$$

$$[K^a_b, R^{c_1 \dots c_6}] = \delta_b^{c_1} R^{ac_2 \dots c_6} + \dots, \quad [K^a_b, R^{c_1 \dots c_3}] = \delta_b^{c_1} R^{ac_2 c_3} + \dots, \quad (A.2)$$

$$[K^a_b, R^{c_1 \dots c_8, d}] = (\delta_b^{c_1} R^{ac_2 \dots c_8, d} + \dots) + \delta_b^d R^{c_1 \dots c_8, a}. \quad (A.3)$$

and

$$[R^{c_1 \dots c_3}, R^{c_4 \dots c_6}] = 2R^{c_1 \dots c_6}, \quad [R^{a_1 \dots a_6}, R^{b_1 \dots b_3}] = 3R^{a_1 \dots a_6 [b_1 b_2, b_3]}, \quad (A.4)$$

where $+\dots$ means the appropriate anti-symmetrisation.

The E_{11} level zero and negative level generators up to level minus three obey the relations

$$[K^a_b, R_{c_1 \dots c_3}] = -\delta_{c_1}^a R_{bc_2 c_3} - \dots, \quad [K^a_b, R_{c_1 \dots c_6}] = -\delta_{c_1}^a R_{bc_2 \dots c_6} - \dots, \quad (A.5)$$

$$[K^a_b, R_{c_1 \dots c_8, d}] = -(\delta_{c_1}^a R_{bc_2 \dots c_8, d} + \dots) - \delta_d^a R_{c_1 \dots c_8, b}. \quad (A.6)$$

$$[R_{c_1 \dots c_3}, R_{c_4 \dots c_6}] = 2R_{c_1 \dots c_6}, \quad [R_{a_1 \dots a_6}, R_{b_1 \dots b_3}] = 3R_{a_1 \dots a_6 [b_1 b_2, b_3]}, \quad (A.7)$$

Finally, the commutation relations between the positive and negative generators are given by

$$\begin{aligned} [R^{a_1 \dots a_3}, R_{b_1 \dots b_3}] &= 18\delta_{[b_1 b_2}^{[a_1 a_2} K^{a_3]}_{b_3]} - 2\delta_{b_1 b_2 b_3}^{a_1 a_2 a_3} D, \quad [R_{b_1 \dots b_3}, R^{a_1 \dots a_6}] = \frac{5!}{2}\delta_{b_1 b_2 b_3}^{[a_1 a_2 a_3} R^{a_4 a_5 a_6]} \\ [R^{a_1 \dots a_6}, R_{b_1 \dots b_6}] &= -5!.3.3\delta_{[b_1 \dots b_5}^{[a_1 \dots a_5} K^{a_6]}_{b_6]} + 5!\delta_{b_1 \dots b_6}^{a_1 \dots a_6} D, \\ [R_{a_1 \dots a_3}, R^{b_1 \dots b_8, c}] &= 8.7.2(\delta_{[a_1 a_2 a_3}^{[b_1 b_2 b_3} R^{b_4 \dots b_8]c} - \delta_{[a_1 a_2 a_3}^{[b_1 b_2 | c]} R^{b_3 \dots b_8]}) \\ [R_{a_1 \dots a_6}, R^{b_1 \dots b_8, c}] &= \frac{7!.2}{3}(\delta_{[a_1 \dots a_6}^{[b_1 \dots b_6} R^{b_7 b_8]c} - \delta_{[a_1 \dots a_6}^{c[b_1 \dots b_5} R^{b_6 b_7 b_8]}) \end{aligned} \quad (A.8)$$

where $D = \sum_b K^b_b$, $\delta_{b_1 b_2}^{a_1 a_2} = \frac{1}{2}(\delta_{b_1}^{a_1} \delta_{b_2}^{a_2} - \delta_{b_1}^{a_2} \delta_{b_2}^{a_1}) = \delta_{b_1}^{[a_1} \delta_{b_2}^{a_2]}$ with similar formulae when more indices are involved. We have taken the liberty of listing the algebra to a somewhat

higher than required in this paper. In fact these equations correct the coefficients of one of the equations contained in reference [3].

The IIA theory arises when we consider the deletion of node ten in the E_{11} Dynkin diagram to leave a $SO(10,10)$ algebra leading to a decomposition of E_{11} into representations of this algebra. However, it is illuminating to then delete node eleven and analyse the representations of $SO(10,10)$ in terms of $SL(10)$. Thus in this case we have two levels which we may write as (l_1, l_2) and are associated with nodes ten and eleven respectively. At level $l_1 = 0$ we find generators with $l_2 = 0$ which are the generators of $GL(10)$, denoted by K^a_b , $a, b = 1, 2 \dots, 10$, and the generator of $\tilde{R}$ which is some combination of $\sum_{a=1}^{11} K^a_a$ and K^{11}_{11} as well as the generators $R^{ab11} \equiv R^{ab}$ and $R_{ab} \equiv R_{ab11}$ at levels $l_2 = 1$ and $l_2 = -1$ respectively. Substituting into the above algebra of equations (A.1-A.8) one finds that these generators obey the algebra

$$\begin{aligned} [K^a_b, K^c_d] &= \delta_b^c K^a_d - \delta_d^a K^c_b, \quad [K^a_b, R^{cd}] = \delta_b^c R^{ad} - \delta_b^d R^{ac}, \quad [K^a_b, R_{cd}] = -\delta_c^a \tilde{R}_{bd} + \delta_d^a \tilde{R}_{bc}, \\ [R^{ab}, R_{cd}] &= 4\delta_{[c}^{[a} K^{b]}_{d]} + \frac{2}{3} \tilde{R}, \quad [R^{ab}, R^{cd}] = 0 = [\tilde{R}_{ab}, \tilde{R}_{cd}], \quad [\tilde{R}, R^{ab}] = 0, \quad [\tilde{R}, R_{ab}] = 0 \end{aligned} \quad (A.9)$$

where $\tilde{R} = -\sum_{a=1}^{11} K^a_a + 3K^{11}_{11}$.

At first sight this is not obviously the algebra of $SO(10,10) \otimes GL(1)$, however, if we redefine the $GL(10)$ generators by

$$\tilde{K}^a_b = K^a_b + \frac{1}{6} \delta_a^b \tilde{R}, \quad (A.10)$$

leaving all other generators the same one, finds the above algebra becomes

$$\begin{aligned} [\tilde{K}^a_b, \tilde{K}^c_d] &= \delta_b^c \tilde{K}^a_d - \delta_d^a \tilde{K}^c_b, \quad [\tilde{K}^a_b, R^{cd}] = \delta_b^c R^{ad} - \delta_b^d R^{ac}, \quad [\tilde{K}^a_b, R_{cd}] = -\delta_c^a \tilde{R}_{bd} + \delta_d^a \tilde{R}_{bc}, \\ [R^{ab}, R_{cd}] &= 4\delta_{[c}^{[a} \tilde{K}^{b]}_{d]}, \quad [R^{ab}, R^{cd}] = 0 = [\tilde{R}_{ab}, \tilde{R}_{cd}] \end{aligned} \quad (A.11)$$

which we recognise as the algebra of $SO(10,10)$ and an additional generator $\tilde{R}$ that commutes with all the $SO(10,10)$ generators.

We also need the fundamental representation of E_{11} associated with node one. By definition this is the representation with highest weight Λ_1 which obeys $(\Lambda_1, \alpha_a) = \delta_{a,1}$, $a = 1, 2 \dots, 11$ where α_a are the simple roots of E_{11} . In the decomposition to $Sl(11)$, corresponding to the deletion of node eleven, one finds that the l_1 representation contains the objects P_a , $a = 1, \dots, 11$, Z^{ab} and $Z^{a_1 \dots a_5}$ corresponding to levels zero, one and two respectively. We have taken the first object, i.e. P_a , to have level zero by choice. Taking these to be generators in a semi-direct product group denoted $E_{11} \otimes_s l_1$ their commutation relation with the level one generators of E_{11} are given by

$$\begin{aligned} [R^{a_1 a_2 a_3}, P_b] &= 3\delta_b^{[a_1} Z^{a_2 a_3]}, \quad [R^{a_1 a_2 a_3}, Z^{b_1 b_2}] = Z^{a_1 a_2 a_3 b_1 b_2}, \\ [R^{a_1 a_2 a_3}, Z^{b_1 \dots b_5}] &= Z^{b_1 \dots b_5 [a_1 a_2, a_3]} + Z^{b_1 \dots b_5 a_1 a_2 a_3} \end{aligned} \quad (A.12)$$

These equations define the normalisation of the generators of the l_1 representation. The commutator of K^a_b with P_c can only take the form $[K^a_b, P_c] = -\delta^a_c P_b + e\delta^a_b P_c$ where e is a constant. This commutator was found [3] to have $e = \frac{1}{2}$ as a result of the fact that l_1 is an E_{11} representation. We summarise the argument here as this relation plays a crucial role in this paper. Since P_1 corresponds to the highest weight state in the representation the action of the Chevalley generator H_{11} is given by $[H_{11}, P_1] = (\alpha_{11}, \Lambda_1)P_1$ where $H_{11} = \frac{2}{3}(K^9_9 + K^{10}_{10} + K^{11}_{11}) - \frac{1}{3}(K^1_1 + \dots + K^8_8)$ is the Chevalley generator in the Cartan subalgebra associated with node eleven. Since $(\Lambda_1, \alpha_{11}) = 0$ we must have $[H_{11}, P_1] = 0$ and then we find $e = \frac{1}{2}$ as claimed. Using the Jacobi identity and the fact that $e = \frac{1}{2}$ we conclude that

$$\begin{aligned} [K^a_b, P_c] &= -\delta^a_c P_b + \frac{1}{2}\delta^a_b P_c, \quad [K^a_b, Z^{c_1 c_2}] = 2\delta^{[c_1}_b Z^{a|c_2]} + \frac{1}{2}\delta^a_b Z^{c_1 c_2}, \\ [K^a_b, Z^{c_1 \dots c_5}] &= 5\delta^{[c_1}_b Z^{a|c_2 \dots c_5]} + \frac{1}{2}\delta^a_b Z^{c_1 \dots c_5} \end{aligned} \quad (A.13)$$

Using the Jacobi identities, equations (A.13) and equations (A.1-A.8) we find that

$$[R^{a_1 \dots a_6}, P_b] = -3\delta^{[a_1}_b Z^{\dots a_6]}, \quad [R^{a_1 \dots a_6}, Z^{b_1 b_2}] = Z^{b_1 b_2 [a_1 \dots a_5, a_6]}, \quad (A.14)$$

and the commutators with the negative root generators are given by

$$[R_{a_1 a_2 a_3}, P_b] = 0, \quad [R_{a_1 a_2 a_3}, Z^{b_1 b_2}] = 6\delta^{b_1 b_2}_{[a_1 a_2} P_{a_3]}, \quad [R_{a_1 a_2 a_3}, Z^{b_1 \dots b_5}] = \frac{5!}{2}\delta^{[b_1 b_2 b_3}_{a_1 a_2 a_3} Z^{b_4 b_5]} \quad (A.15)$$

As seen from the IIA perspective we first delete node ten and then node eleven; the level $l_1 = 0$ generators of the l_1 multiplet are P_a , $a = 1, \dots, 10$ and $Q^a \equiv -2Z^{a11}$ which have levels $l_2 = 0$ and $l_2 = 1$ respectively. We find from the above algebra that they obey the commutation relations

$$\begin{aligned} [K^c_b, P_a] &= -\delta^c_a P_b + \frac{1}{2}\delta^c_b P_a, \quad [R^{ab}, P_c] = -\frac{1}{2}(\delta^a_c Q^b - \delta^b_c Q^a) = -\delta^{[a}_c Q^{b]}, \quad [\tilde{R}_{ab}, P_c] = 0, \\ [K^a_b, Q^c] &= \delta^c_b Q^a + \frac{1}{2}\delta^c_b Q^a, \quad [\tilde{R}_{ab}, Q^c] = 2(\delta^c_a P_b - \delta^c_b P_a) = 4\delta^c_{[a} P_{b]}, \quad [R^{ab}, Q^c] = 0, \\ [\tilde{R}, P_a] &= -3P_a, \quad [\tilde{R}, Q^a] = -3Q^a \end{aligned} \quad (A.16)$$

It will be useful to also give the commutators in terms of the generator $R = \frac{1}{12}(-\sum_{a=1}^{11} K^a_a + 9K^{11}_{11})$, used in the non-linear realisation, which leads to

$$[R, P_a] = 0, \quad [R, Q^a] = \frac{1}{2}Q^a, \quad [R, R^{ab}] = \frac{1}{2}R^{ab}, \quad [R, R_{ab}] = -\frac{1}{2}R_{ab} \quad (A.17)$$

It is instructive to give these relations when we take the generators of equation (A.10) that are those of the $SO(10,10)$ algebra. Then one finds that

$$[\tilde{K}^c_b, P_a] = -\delta^c_a P_b, \quad [R^{ab}, P_c] = -\frac{1}{2}(\delta^a_c Q^b - \delta^b_c Q^a) = -\delta^{[a}_c Q^{b]}, \quad [\tilde{R}_{ab}, P_c] = 0,$$

$$[\tilde{K}^a{}_b, Q^c] = \delta_b^c Q^a, [\tilde{R}_{ab}, Q^c] = 2(\delta_a^c P_b - \delta_b^c P_a) = 4\delta_{[a}^c P_{b]}, [R^{ab}, Q^c] = 0, \quad (A.18)$$

which are those corresponding to the usual vector representation of $SO(10,10)$. The relations with the additional $GL(1)$ factor are given in equation (A.16).




Fig 1. The E_{11} Dynkin diagram

References

- [1] P. West, E_{11} and M Theory, *Class. Quant. Grav.* **18** (2001) 4443, [hep-th/0104081](#)
- [2] P. C. West, *Hidden superconformal symmetry in M theory*, *JHEP* **08** (2000) 007, [hep-th/0005270](#)
- [3] P. West, E_{11} , $SL(32)$ and Central Charges, *Phys. Lett. B* **575** (2003) 333-342, [hep-th/0307098](#)
- [4] P. West, E_{11} origin of brane charges and U -duality multiplets, *JHEP* **0408** (2004) 052, [hep-th/0406150](#).
- [5] A. Kleinschmidt and P. West, *Representations of $\mathcal{G}^{+++}$ and the role of space-time*, *JHEP* **0402** (2004) 033, [hep-th/0312247](#).
- [6] P. West, Brane dynamics, central charges and E_{11} , *JHEP* **0503** (2005) 077, [hep-th/0412336](#).
- [7] P. Cook and P. West, Charge multiplets and masses for $E(11)$; *JHEP* **11** (2008) 091, [arXiv:0805.4451](#).
- [8] F. Riccioni and P. West, $E(11)$ -extended space-time and gauged supergravities, *JHEP* **0802**:039,2008; [hep-th/0712.1795](#)
- [9] N. Hitchin, Generalized Calabi-Yau manifolds, *Q. J. Math.* **54** (2003), no. 3, 281, [math.DG/0209099](#); Brackets, form and invariant functionals, [math.DG/0508618](#).
- [10] M. Gualtieri, Generalized complex geometry, PhD Thesis (2004), [math.DG/0401221v1](#).
- [11] Borisov and V. Ogievetsky, Theory of dynamical affine and conformal symmetries as gravity theory of the gravitational field, *Theor. Math. Phys.* **21** (1975) 1179.
- [12] C. Hillmann, Generalized $E(7(7))$ coset dynamics and $D=11$ supergravity, *JHEP* **0903**, 135 (2009), [hep-th/0901.1581](#).
- [13] M. Duff, Duality Rotations In String Theory, *Nucl. Phys. B* **335** (1990) 610; M. Duff and J. Lu, Duality rotations in membrane theory, *Nucl. Phys. B* **347** (1990) 394.
- [14] A. Tseytlin, Duality Symmetric Formulation Of String World Sheet Dynamics, *Phys.Lett. B* **242** (1990) 163, Duality Symmetric Closed String Theory And Interacting Chiral Scalars, *Nucl. Phys. B* **350**, 395 (1991).
- [15] P. West, Generalised space-time and duality, [hep-th/1006.0893](#).
- [16] I. Schnakenberg and P. West, Kac-Moody Symmetries of IIB Supergravity, *Phys.Lett. B* **517** (2001) 421-428, [hep-th/0107181](#).
- [17] C. Hull and B. Zwiebach, Double Field Theory, *JHEP* **0909** (2009) 099, [hep-th/0904.4664](#).

- [18] C. Hull and B. Zwiebach, The gauge algebra of double field theory and Courant brackets, JHEP **0909** (2009) 090, hep-th/0908.1792.
- [19] O. Hohm, C. Hull and B. Zwiebach, Background independent action for double field theory, hep-th/1003.5027.
- [20] O. Hohm, C. Hull and B. Zwiebach, Generalised metric formulation of double field theory, hep-th/1006.4823.
- [21] T. Kugo and B. Zwiebach, Target space duality as a symmetry of string field theory, Prog. Theor. Phys. **87**, 801 (1992) hep-th/9201040.
- [22] W. Siegel, Superspace duality in low-energy superstrings, Phys. Rev. D **48**, 2826 (1993), hep-th/9305073; Two vierbein formalism for string inspired axionic gravity, Phys. Rev. D **47**, 5453 (1993) hep-th/9302036.
- [23] T. Kugo and B. Zweibach, Prog. Theor. Phys. bf 87, (1992) 801.
- [24] Seminar of Olaf Holm given at the Solvay workshop, "Symmetries and dualities in gravitational theories" in Brussels, May 21, 2010.
- [25] Discussion session after the seminar of Olaf Holm specified in the previous reference.